

Stufenschema Analgo-Sedation Neugeborene auf Intensivstation

<div style="border: 1px solid black; padding: 5px; display: inline-block;"> 23 ⁰/₇ - 27 ⁶/₇ Sedation NEC-konservativ </div>				Rocuronium 0.3 mg/kg
				< 3 Tage: Fentanyl DT 0.5-3 mcg/kg/h >3 Tage: ± Dexmedetomidine-DT 0.2-1 mcg/kg/h
Fentanyl Boli 1-2 mcg/kg/Dosis				
Nicht-medikamentöse Therapie + Glucoselösung				
Stufe 1	Stufe 2	Stufe 3	Stufe 4	

<div style="border: 1px solid black; padding: 5px; display: inline-block;"> 23 ⁰/₇ - 27 ⁶/₇ Postoperativ </div>				Rocuronium 0.3 mg/kg
				Fentanyl Boli 1-2 mcg/kg/Dosis >3 Tage: ± Dexmedetomidine 0.2-1 mcg/kg/h
Paracetamol 10mg/kg 12h i.v. + Fentanyl-DT 0.5-3(-5) mcg/kg/h				
Nicht-medikamentöse Therapie + Glucoselösung				
Stufe 1	Stufe 2	Stufe 3	Stufe 4	

<div style="border: 1px solid black; padding: 5px; display: inline-block;"> >28 ⁰/₇ Sedation </div>				Rocuronium 0.3 mg/kg
				28 ⁰ / ₇ - 31 ⁶ / ₇ : Morphin DT 5-15 mcg/kg/h >32 ⁰ / ₇ : Morphin DT 5-20 mcg/kg/h kreislaufinstabil: Fentanyl DT 0.5-3 mcg/kg/h
Morphin Boli 25-50 mcg/kg/Dosis kreislaufinstabil: Fentanyl Boli 1-2 mcg/kg/Dosis				
Nicht-medikamentöse Therapie + Glucoselösung				
Stufe 1	Stufe 2	Stufe 3	Stufe 4	

<div style="border: 1px solid black; padding: 5px; display: inline-block;"> <p>>28 0/7 Postoperativ</p> </div>				Rocuronium 0.3 mg/kg
				<p>Fentanyl Boli 1-2 mcg/kg/Dosis >3 Tage: ± Dexmedetomidine 0.2-1(-1.5) mcg/kg/h</p>
<p>Paracetamol 10mg/kg 6-12h i.v. + Fentanyl-DT 0.5-3(-5) mcg/kg/h</p>				
<p>Nicht-medikamentöse Therapie + Glucoselösung</p>				
Stufe 1	Stufe 2	Stufe 3	Stufe 4	

<div style="border: 1px solid black; padding: 5px; display: inline-block;"> <p>>28 0/7 Pulmonalarterielle Hypertonie Mekoniumaspiration Zwerchfellhernie</p> </div>				Rocuronium 1 mg/kg
				<p>Morphin Boli 25-50 mcg/kg/Dosis kreislaufinstabil: Fentanyl Boli 1-2 mcg/kg/Dosis >3 Tage: Dexmedetomidine 0.2-1(-1.5) mcg/kg/h</p>
<p>Morphin Bolus 50mcg/kg + Morphin DT 5-15 mcg/kg/h kreislaufinstabil: Fentanyl Bolus + Fentanyl DT 0.5-3 mcg/kg/h</p>				
<p>Nicht-medikamentöse Therapie + Glucoselösung</p>				
Stufe 1	Stufe 2	Stufe 3	Stufe 4	

<div style="border: 1px solid black; padding: 5px; display: inline-block;"> <p>>35 0/7 ther. Hypothermie</p> </div>				Rocuronium 0.3 mg/kg
				<p>Morphin DT 10 mcg/kg/h kreislaufinstabil: Fentanyl DT 0.5-3 mcg/kg/h</p>
<p>Morphin Boli 25-50mcg/kg/Dosis kreislaufinstabil: Fentanyl-Boli 1-2 mcg/kg/Dosis</p>				
<p>Nicht-medikamentöse Therapie + Glucoselösung</p>				
Stufe 1	Stufe 2	Stufe 3	Stufe 4	